

Physics at Hadron Colliders

Lecture II

Beate Heinemann

*University of California, Berkeley and
Lawrence Berkeley National Laboratory*

CERN, Summer Student Lectures, 2010

Outline

- Lecture I: Introduction
 - Outstanding problems in particle physics
 - and the role of hadron colliders
 - Current colliders: Tevatron and LHC
 - Hadron-hadron collisions
- Lecture II: Standard Model Measurements
 - Standard Model Cross Section Measurements as Tests of QCD
 - Precision measurements in electroweak sector
- Lecture III: Searches for the Higgs Boson
 - Standard Model Higgs Boson
 - Higgs Bosons beyond the Standard Model
- Lecture IV: Searches for New Physics
 - Supersymmetry
 - High Mass Resonances (Extra Dimensions etc.)

Standard Model Cross Section Measurements as test of QCD

- **Jets**
- **W and Z bosons**
- **Top Quark Production**


What is a Cross Section?

- Differential cross section: $d\sigma/d\Omega$:
 - Probability of a scattered particle in a given quantum state per solid angle $d\Omega$
 - E.g. Rutherford scattering experiment
- Other differential cross sections: $d\sigma/dE_T(\text{jet})$
 - Probability of a jet with given E_T
- Integrated cross section
 - Integral: $\sigma = \int d\sigma/d\Omega d\Omega$



Measurement:

$$\sigma = (N_{\text{obs}} - N_{\text{bg}}) / (\epsilon L)$$

Jet Cross Sections




- Inclusive jets: processes qq , qg , gg




- Highest E_T probes shortest distances
 - Tevatron: $r_q < 10^{-18}$ m
 - LHC: $r_q < 10^{-19}$ m (?)
 - Could e.g. reveal substructure of quarks
- Tests perturbative QCD at highest energies

Jet Cross Section History




- Run I (1996):
 - Excess at high E_T
 - Could be signal for quark substructure?!?



Jet Cross Section History



Inclusive Jet cross section




■ Since Run I:

■ Revision of parton density functions


- Gluon is uncertain at high x
- It including these data describes data well






Jet Cross Sections: Tevatron



- Excellent agreement with QCD calculation over 8 orders of magnitude!
- No excess any more at high E_T
 - Large pdf uncertainties will be constrained by these data




Jet Cross Sections: Tevatron and LHC



■ First LHC measurements agree well with theoretical predictions

High Mass Dijet Event: $M=1.4$ TeV




CDF Run II Preliminary

Jet E_{T1} = 666 GeV (corr)
583 GeV (raw)

η_{1} = 0.31 (detector)
0.43 (corr z)

Jet E_{T2} = 633 GeV (corr)
546 GeV (raw)

η_{2} = -0.30 (detector)
-0.19 (corr z)






Run 152507
Event 1222318

DiJet Mass = 1364 GeV (corr)

z vertex = -25 cm

High-Mass Di-Jets at the LHC

- $M(jj)=2.55$ GeV
- $pT(j1)=420$ GeV
- $pT(j2)=320$ GeV





 **ATLAS**
EXPERIMENT

Run Number: 158548, Event Number: 5917927

Date: 2010-07-04 07:24:40 CEST

W and Z Bosons


- Focus on leptonic decays:
 - Hadronic decays ~impossible due to enormous QCD dijet background
- Selection:
 - Z:
 - Two leptons $p_T > 20$ GeV
 - Electron, muon, tau
 - W:
 - One lepton $p_T > 20$ GeV
 - Large imbalance in transverse momentum
 - Missing $E_T > 20$ GeV
 - Signature of undetected particle (neutrino)
- Excellent calibration signal for many purposes:
 - Electron energy scale
 - Track momentum scale
 - Lepton ID and trigger efficiencies
 - Missing E_T resolution
 - Luminosity ...



Lepton Identification


■ Electrons:

- compact electromagnetic cluster in calorimeter
- Matched to track





■ Muons:

- Track in the muon chambers
- Matched to track




■ Taus:

- Narrow jet
- Matched to one or three tracks




■ Neutrinos:

- Imbalance in transverse momentum
- Inferred from total transverse energy measured in detector
- More on this in Lecture 4




Electron and Muon Identification

- Desire:
 - High efficiency for isolated electrons
 - Low misidentification of jets




- Performance:
 - Efficiency:
 - 60-100% depending on $|\eta|$
 - Measured using Z's



Electrons and Jets

Run 166892 Evt 2775140 Sun Oct 27 03:15:49 2002


ET scale: 49 GeV



Run 166892 Evt 3223863 Sun Oct 27 03:43:08 2002

E scale: 20 GeV

Hadronic Calorimeter Energy




+2

Electromagnetic Calorimeter Energy

- Jets can look like electrons, e.g.:
 - photon conversions from π^0 's: ~13% of photons convert (in CDF)
 - early showering charged pions
- And there are lots of jets!!!


Jets faking Electrons

- Jets can pass electron ID cuts,
 - Mostly due to
 - early showering charged pions
 - Conversions: $\pi^0 \rightarrow \gamma\gamma \rightarrow ee + X$
 - Semileptonic b-decays
 - Difficult to model in MC
 - Hard fragmentation
 - Detailed simulation of calorimeter and tracking volume
- Measured in inclusive jet data at various E_T thresholds
 - Prompt electron content negligible:
 - $N_{\text{jet}} \sim 10$ billion at 50 GeV!
 - Fake rate per jet:
 - CDF, tight cuts: 1/10000
 - ATLAS, tight cuts: 1/80000
 - Typical uncertainties 50%



Jets faking “loose” electrons

Fake Rate (%)



W's and Z's

Z mass reconstruction

- Invariant mass of two leptons

$$m = \sqrt{(E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2}$$



- Sets electron energy scale by comparison to LEP measured value

W mass reconstruction

- Do not know neutrino p_z
- No full mass reconstruction possible
- Transverse mass:


$$m_T = \sqrt{|p_T^\ell|^2 + |p_T^\nu|^2 - (\vec{p}_T^\ell + \vec{p}_T^\nu)^2}$$

Di-Electron Invariant Mass Spectrum





Tevatron W and Z Cross Section Results

- Uncertainties:
 - Experimental: 2%
 - Theoretical: 2%
 - Luminosity: 6%
- Can we use these processes to normalize luminosity?
 - Is theory reliable enough?




LHC signals of W's and Z's with $\sim 0.3 \text{ pb}^{-1}$



- $0.2\text{-}0.3 \text{ pb}^{-1}$ yield clean signals of W's and Z's


LHC W and Z Cross Sections



- Data agree well with theoretical expectation
 - Uncertainties: 13% (W), 17% (Z)
 - Luminosity uncertainty 10%

Top Quark Production and Decay

- At Tevatron, mainly produced in pairs via the strong interaction




- Decay via the electroweak interactions $\text{Br}(t \rightarrow Wb) \sim 100\%$
Final state is characterized by the decay of the W boson

Dilepton

Lepton+Jets

All-Jets




Different sensitivity and challenges in each channel

How to identify the top quark

SM: $t\bar{t}$ pair production, $\text{Br}(t \rightarrow bW) = 100\%$, $\text{Br}(W \rightarrow lv) = 1/9 = 11\%$

dilepton	(4/81)	2 leptons + 2 jets + missing E_T
l+jets	(24/81)	1 lepton + 4 jets + missing E_T
fully hadronic	(36/81)	6 jets


(here: $l = e, \mu$)



How to identify the top quark

SM: $t\bar{t}$ pair production, $\text{Br}(t \rightarrow bW) = 100\%$, $\text{Br}(W \rightarrow lv) = 1/9 = 11\%$


dilepton	(4/81)	2 leptons + 2 jets + missing E_T
lepton+jets	(24/81)	1 lepton + 4 jets + missing E_T
fully hadronic	(36/81)	6 jets



How to identify the top quark

SM: $t\bar{t}$ pair production, $\text{Br}(t \rightarrow bW) = 100\%$, $\text{Br}(W \rightarrow lv) = 1/9 = 11\%$


dilepton	(4/81)	2 leptons + 2 jets + missing E_T
lepton+jets	(24/81)	1 lepton + 4 jets + missing E_T
fully hadronic	(36/81)	6 jets




How to identify the top quark

SM: $t\bar{t}$ pair production, $\text{Br}(t \rightarrow bW) = 100\%$, $\text{Br}(W \rightarrow lv) = 1/9 = 11\%$

dilepton	(4/81)	2 leptons + 2 jets + missing E_T
lepton+jets	(24/81)	1 lepton + 4 jets + missing E_T
fully hadronic	(36/81)	6 jets




Top Event Categories




Finding the Top at Tevatron and LHC


- **B-tagging helps a lot:**
 - Signal/Background improved by about a factor of 10 when using b-tagging
- **Tevatron (with 4 jets):**
 - no b-tagging: $S/B \approx 0.8$
 - With b-tagging: $S/B \approx 6$
- **Use b-tagging both at Tevatron and at LHC**



Finding the b-jets






- Exploit large lifetime of the b-hadron
 - B-hadron flies before it decays: $d=c\tau$
 - Lifetime $\tau = 1.5 \text{ ps}^{-1}$
 - $d=c\tau = 460 \text{ }\mu\text{m}$
 - Can be resolved with silicon detector resolution
- Procedure “Secondary Vertex”:
 - reconstruct primary vertex:
 - resolution $\sim 30 \text{ }\mu\text{m}$
 - Search tracks inconsistent with primary vertex (large d_0):
 - Candidates for secondary vertex
 - See whether three or two of those intersect at one point
 - Require displacement of secondary from primary vertex
 - Form L_{xy} : transverse decay distance projected onto jet axis:
 - $L_{xy} > 0$: b-tag along the jet direction => real b-tag or mistag
 - $L_{xy} < 0$: b-tag opposite to jet direction => mistag!
 - Significance: e.g. $\delta L_{xy} / L_{xy} > 7$ (i.e. 7σ significant displacement)
- More sophisticated techniques exist



The Top Signal: Lepton + Jets

- Select:
 - 1 electron or muon
 - Large missing E_T
 - 1 or 2 b-tagged jets



double-tagged events, nearly no background




Check backgrounds

Top Signal

$$\sigma(t\bar{t}) = 8.3^{+0.6}_{-0.5}(\text{stat}) \pm 1.1(\text{syst}) \text{ pb}$$


The Top Signal: Dilepton

- Select:
 - 2 leptons: $ee, e\mu, \mu\mu$
 - Large missing E_T
 - 2 jets (with or w/o b-tag)




$$\sigma = 6.2 \pm 0.9 \text{ (stat)} \pm 0.9 \text{ (sys)} \text{ pb}$$

The Top Cross Section



- Many measurements that use different techniques
- Good agreement
 - between all measurements
 - between data and theory
- Experimental Precision: $\sim 6\%$
 - Better than theoretical precision

Searching for Top at CMS and ATLAS





- Both experiments have seen a few candidate events
- No conclusive signal yet

Precision Measurement of Electroweak Sector of the Standard Model

- **W boson mass**
- **Top quark mass**
- **Implications for the Higgs boson**



The W boson, the top quark and the Higgs boson

- Top quark is the heaviest known fundamental particle
 - Today: $m_{\text{top}} = 173.3 \pm 1.1$ GeV
 - Run 1: $m_{\text{top}} = 178 \pm 4.3$ GeV/c²
- Masses related through radiative corrections:
 - $m_W \sim M_{\text{top}}^2$
 - $m_W \sim \ln(m_H)$
- If there are new particles the relation might change:
 - Precision measurement of top quark and W boson mass can reveal new physics





The W boson, the top quark and the Higgs boson

- Top quark is the heaviest known fundamental particle
 - Today: $m_{\text{top}} = 173.3 \pm 1.1$ GeV
 - Run 1: $m_{\text{top}} = 178 \pm 4.3$ GeV/c²
- Masses related through radiative corrections:
 - $m_W \sim M_{\text{top}}^2$
 - $m_W \sim \ln(m_H)$
- If there are new particles the relation might change:
 - Precision measurement of top quark and W boson mass can reveal new physics




The W boson, the top quark and the Higgs boson

- Top quark is the heaviest known fundamental particle
 - Today: $m_{\text{top}} = 173.3 \pm 1.1$ GeV
 - Run 1: $m_{\text{top}} = 178 \pm 4.3$ GeV/c²
- Masses related through radiative corrections:
 - $m_W \sim M_{\text{top}}^2$
 - $m_W \sim \ln(m_H)$
- If there are new particles the relation might change:
 - Precision measurement of top quark and W boson mass can reveal new physics



W Boson mass

- Real **precision** measurement:
 - LEP: $M_W = 80.367 \pm 0.033 \text{ GeV}/c^2$
 - Precision: 0.04%
 - => Very challenging!
- Main measurement ingredients:
 - **Lepton p_T**
 - **Hadronic recoil** parallel to lepton: $u_{||}$
- $Z \rightarrow ll$ superb calibration sample:
 - but statistically limited:
 - About a factor 10 less Z's than W's
 - Most systematic uncertainties are related to size of Z sample
 - Will scale with $1/\sqrt{N_Z}$ ($=1/\sqrt{L}$)







$$m_T = \sqrt{2p_T^l \cancel{p}_T (1 - \cos \Delta\phi)},$$

$$\cancel{p}_T \approx |p_T + u_{||}|$$

$$m_T \approx 2p_T \sqrt{1 + u_{||}/p_T} \approx 2p_T + u_{||}$$

Lepton Momentum Scale and Resolution



- Systematic uncertainty on momentum scale: 0.04%

Systematic Uncertainties

m_T Fit Uncertainties			
Source	$W \rightarrow \mu\nu$	$W \rightarrow e\nu$	Correlation
Tracker Momentum Scale	17	17	100%
Calorimeter Energy Scale	0	25	0%
Lepton Resolution	3	9	0%
Lepton Efficiency	1	3	0%
Lepton Tower Removal	5	8	100%
Recoil Scale	9	9	100%
Recoil Resolution	7	7	100%
Backgrounds	9	8	0%
PDFs	11	11	100%
W Boson p_T	3	3	100%
Photon Radiation	12	11	100%
Statistical	54	48	0%
Total	60	62	-



Limited by data statistics

Limited by data and theoretical understanding

TABLE IX: Uncertainties in units of MeV on the transverse mass fit for m_W in the $W \rightarrow \mu\nu$ and $W \rightarrow e\nu$ samples.

- Overall uncertainty 60 MeV for both analyses
 - Careful treatment of correlations between them
- Dominated by stat. error (50 MeV) vs syst. (33 MeV)

W Boson Mass



World average:

$$M_W = 80399 \pm 23 \text{ MeV}$$


Ultimate precision:

Tevatron: 15-20 MeV

LHC: unclear (5 MeV?)

Top Mass Measurement: $t\bar{t} \rightarrow (bl\nu)(bqq)$

- 4 jets, 1 lepton and missing E_T
 - Which jet belongs to what?
 - Combinatorics!
- B-tagging helps:
 - 2 b-tags \Rightarrow 2 combinations
 - 1 b-tag \Rightarrow 6 combinations
 - 0 b-tags \Rightarrow 12 combinations
- Two Strategies:
 - Template method:
 - Uses “best” combination
 - Chi2 fit requires $m(t) = m(\bar{t})$
 - Matrix Element method:
 - Uses all combinations
 - Assign probability depending on kinematic consistency with top



Top Mass Determination

- Inputs:

- Jet 4-vectors
- Lepton 4-vector
- Remaining transverse energy, $p_{T,UE}$:

- $p_{T,v} = -(p_{T,l} + p_{T,UE} + \sum p_{T,jet})$

- Constraints:





- $M(l\nu) = M_W$
- $M(q\bar{q}) = M_W$
- $M(t) = M(\bar{t})$

- Unknown:

- Neutrino p_z

- 1 unknown, 3 constraints:


- Overconstrained
- Can measure $M(t)$ for each event: m_t^{reco}
- Leave jet energy scale (“JES”) as free parameter



Selecting correct combination
20-50% of the time

Example Results on m_{top}

CDF Run II Preliminary 5.6 fb^{-1}




$$m_{\text{top}} = 173.7 \pm 0.8 \pm 1.6 \text{ GeV} \quad (L=2.6 \text{ fb}^{-1})$$



$$m_{\text{top}} = 173.0 \pm 0.6 \pm 1.1 \text{ GeV} \quad (L=5.6 \text{ fb}^{-1})$$

Combining M_{top} Results

- Excellent results in each channel
 - Dilepton
 - Lepton+jets
 - All-hadronic
- Combine them to improve precision
 - Include Run-I results
 - Account for correlations
- **Uncertainty: 1.1 GeV (0.61%)**
 - Dominated by syst. uncertainties
- Precision so high that theorists wonder about what it's exact definition is!





Implications for the Higgs Boson

LEPEWWG July 2010

Relation: M_W vs m_{top} vs M_H

$$m_H = 89^{+35}_{-26} \text{ GeV}$$



Standard Model still works!

Indirect constraints:
 $m_H < 158 \text{ GeV @95\%CL}$

Conclusions

- Perturbative QCD describes hadron collider data successfully:
 - Jet cross sections: $\Delta\sigma/\sigma \approx 20\text{-}100\%$ (at Tevatron and LHC)
 - W/Z cross section: $\Delta\sigma/\sigma \approx 6\%$ (Tevatron) and $\sim 15\%$ (LHC)
 - Top cross section: $\Delta\sigma/\sigma \approx 6\%$
 - Starting to probe top at the LHC
- High Precision measurements at Tevatron
 - W boson mass: $\Delta M_W/M_W = 0.028\%$
 - top quark mass: $\Delta m_{\text{top}}/m_{\text{top}} = 0.61\%$
- Standard Model still works!
 - Higgs boson constrained
 - $114 < m_H < 158 \text{ GeV}/c^2$ at 95% C.L.
 - Direct Searches: see next lecture!

Luminosity Measurement

$$R_{pp} = \mu_{pp} \cdot f_{BC} = \sigma_{inel} \cdot \epsilon_{pp} \cdot \delta(L) \cdot L$$

L - luminosity

f_{bc} - Bunch Crossing rate

μ_a - # of pp / BC

σ_{LM}

σ_{inel} - inelastic x-section

ϵ_{pp} - acceptance for a single pp

$\delta(L)$ - detector non-linearity

- Measure events with 0 interactions
 - Related to R_{pp}
- Normalize to measured inelastic pp cross section
 - Tevatron: 60.7 ± 2.4 mb
- LHC: 70-120 mb

